

# Bayer plc



Bayer House  
Strawberry Hill  
Newbury  
Berkshire RG14 1JA  
United Kingdom

Telephone: +44 (0)1635 563 000  
Facsimile: +44 (0)1635 563 393  
Company Web Site: <http://www.bayer.co.uk>

Due to regulatory changes, the content of the following Patient Information Leaflet may vary from the one found in your medicine pack. Please compare the 'Leaflet prepared/revised date' towards the end of the leaflet to establish if there have been any changes.

If you have any doubts or queries about your medication, please contact your doctor or pharmacist.



## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Levitra 10 mg orodispersible tablets Vardenafil

#### Read all of this leaflet carefully before you start using this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### In this leaflet:

1. What Levitra is and what it is used for
2. Before you take Levitra
3. How to take Levitra
4. Possible side effects
5. How to store Levitra
6. Further information

## 1. WHAT LEVITRA IS AND WHAT IT IS USED FOR

Levitra contains vardenafil, a member of a class of medicines called phosphodiesterase type 5 inhibitors. They are used for the treatment of erectile dysfunction in adult men, a condition which implies difficulties in getting or keeping an erection.

At least one in ten men has trouble getting or keeping an erection at some time. There may be physical or psychological causes, or a mixture of both. Whatever the cause is, due to muscle and blood vessel changes not enough blood stays in the penis to make it hard and keep it hard.

Levitra will only work when you are sexually stimulated. It reduces the action of the natural chemical in your body which makes erections go away. Levitra allows an erection to last long enough for you to satisfactorily complete sexual activity.

## 2. BEFORE YOU TAKE LEVITRA

### Do not take Levitra

- If you are allergic (hypersensitive) to vardenafil or any of the other ingredients of Levitra. See the ingredients in section 6. Signs of an allergic reaction include a rash, itching, swollen face or lips and shortness of breath.
- If you are taking medicines containing nitrates, such as glycerol trinitrate for angina, or nitric oxide donors, such as amyl nitrite. Taking these medicines with Levitra could seriously affect your blood pressure.
- If you are taking ritonavir or indinavir, medicines used to treat human immunodeficiency virus (HIV) infections.
- If you are over 75 years of age and are taking ketoconazole or itraconazole, anti-fungal medicines
- If you have a severe heart or liver problem
- If you are having kidney dialysis
- If you have recently had a stroke or heart attack
- If you have or have had low blood pressure
- If your family has a history of degenerative eye diseases (such as *retinitis pigmentosa*)
- If you have ever had a condition involving loss of vision due to damage to the optic nerve from insufficient blood supply known as non-arteritic ischemic optic neuropathy (NAION).

### Take special care with Levitra

- If you have heart trouble. It may be risky for you to have sex
- If you suffer from irregular heart beat (cardiac arrhythmia) or inherited heart diseases affecting your electrocardiogram
- If you have a physical condition affecting the shape of the penis. This includes conditions called angulation, Peyronie's disease and cavernosal fibrosis
- If you have an illness that can cause erections which won't go away (priapism). These include sickle cell disease, multiple myeloma and leukaemia
- If you have stomach ulcers (also called gastric or peptic ulcers)
- If you have a bleeding disorder (such as haemophilia)
- If you are using any other treatments for erection difficulties, including Levitra film-coated tablets. See "Using other medicines".
- If you experience sudden decrease or loss of vision, stop taking Levitra and contact your doctor immediately.

### Children

Levitra is not intended for use by children or adolescents under 18.

### **Using other medicines**

Please tell your doctor or pharmacist if you are using or have recently used any other medicines, including medicines obtained without a prescription.

Some medicines may cause problems, especially these:

- Nitrates, medicines for angina, or nitric oxide donors, such as amyl nitrite. Taking these medicines with Levitra could seriously affect your blood pressure. Talk to a doctor without taking Levitra
- Medicine for the treatment of arrhythmias, such as quinidine, procainamide, amiodarone or sotalol
- Ritonavir or indinavir, medicines for HIV. Talk to a doctor without taking Levitra
- Ketoconazole or itraconazole, anti-fungal medicines
- Erythromycin, or clarithromycin, macrolide antibiotics
- Alpha-blockers, a type of medicine used to treat high blood pressure and enlargement of the prostate (as benign prostatic hyperplasia).

Do not use Levitra orodispersible tablets combined with any other treatment for erectile dysfunction, including Levitra film-coated tablets.

### **Taking Levitra with food and drink**

- You can take Levitra orodispersible tablets with or without food, but do not take this medicine with any liquid.
- Don't drink grapefruit juice when you use Levitra. It can interfere with the usual effect of the medicine.
- Alcoholic drink can make erection difficulties worse.

### **Pregnancy and breast-feeding**

Levitra is not for use by women.

### **Driving and using machines**

Levitra might make some people feel dizzy or affect their vision. If you feel dizzy, or if your vision is affected after taking Levitra don't drive or operate any tools or machines.

### **Important information about some of the ingredients of Levitra 10 mg orodispersible tablets:**


- Aspartame is a source of phenylalanine, and may be harmful for people with a disorder called phenylketonuria.
- Sorbitol: If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

## **3. HOW TO TAKE LEVITRA ORODISPERSIBLE TABLETS**

Always take Levitra exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. The usual dose is 10 mg.

Take Levitra about 60 minutes before sexual activity. With sexual stimulation you may achieve an erection anywhere from 25 minutes up to four to five hours after taking Levitra.

- Do not remove the orodispersible tablet from the blister until you are going to take it. With dry hands, pull the blister edge from the blister holder and press gently on top to release the tablet on your hand. Do not crush the tablet.
- Place the entire orodispersible tablet in the mouth, on the tongue, where it will dissolve in seconds, then swallow with saliva. The orodispersible tablet must be taken without any liquid.



**Do not take Levitra orodispersible tablets** with any other forms of Levitra.

**Do not take Levitra** more than once a day.

Tell your doctor if you think Levitra is too strong or too weak. He or she may suggest a switch to an alternative Levitra formulation with a different dose, depending on how well it works for you.

**If you take more Levitra than you should**

Men who take too much Levitra may experience more side effects or may get severe back pain. If you take more Levitra than you should, tell your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### **4. POSSIBLE SIDE EFFECTS**

Like all medicines, Levitra can cause side effects, although not everybody gets them. Most of the effects are mild or moderate.

Partial, sudden, temporary or permanent decrease or loss of vision in one or both eyes has been experienced by patients. Stop taking Levitra and contact your doctor immediately.  
Sudden decrease or loss of hearing has been reported.

The chance of having a side effect is described by the following categories:

**Very common** (*affects more than 1 user in 10*)

- Headache

**Common** (*affects 1 to 10 users in 100*)

- Dizziness
- Flushing
- Blocked or runny nose
- Indigestion

**Uncommon** (*affects 1 to 10 users in 1,000*)

- Swelling of skin and mucous tissue including swollen face, lips or throat.
- Sleep disorder
- Numbness and impaired perception of touch
- Sleepiness
- Effects on vision; redness of the eye, effects on colour vision, eye pain and discomfort, light sensitivity
- Ringing in the ears; vertigo
- Fast heart beat or pounding heart
- Breathlessness
- Stuffy nose
- Acid reflux, gastritis, abdominal pain, diarrhoea, vomiting; feeling sick (nausea), dry mouth
- Raised levels of liver enzymes in your blood

- Rash, reddened skin
- Back or muscle pain; increase in blood of a muscle enzyme (creatine phosphokinase), muscle stiffness
- Prolonged erections
- Malaise

**Rare** (*affects 1 to 10 users in 10,000*)

- Inflammation of the eyes (conjunctivitis)
- Allergic reaction
- Anxiety
- Fainting
- Amnesia
- Seizure
- Increased pressure in the eye (glaucoma), lacrimation increased
- Effects on the heart (such as heart attack, altered heart beat or angina)
- High or low blood pressure
- Nose bleed
- Effect on results of blood tests to check liver function.
- Sensitivity of the skin to sun light
- Painful erections
- Chest pain

If any of the side effects gets serious or if you notice any side effects not listed in this leaflet, please tell your doctor.

## 5. HOW TO STORE LEVITRA

Keep out of the reach and sight of children.

Do not use Levitra after the expiry date which is stated on the carton after “EXP”. The expiry date refers to the last day of that month.

Store in the original package in order to protect from humidity and light.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## 6. FURTHER INFORMATION

### What Levitra orodispersible tablets contains

- The active substance is vardenafil. Each orodispersible tablet contains 10 mg of vardenafil (as hydrochloride).
- The other ingredients of the tablets are: Magnesium stearate, aspartame (E951), peppermint flavour, mannitol (E421) sorbitol (E420), crospovidone and silica colloidal hydrated.

### What Levitra 10 mg orodispersible tablets looks like and contents of the pack

Levitra 10 mg orodispersible tablets are round and white. They are provided in blister packs containing 1, 2 or 4 tablets.

Not all pack sizes may be marketed.

## Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder  
Bayer Schering Pharma AG,  
13342 Berlin,  
Germany.

Manufacturer  
Bayer Schering Pharma AG,  
D-51368 Leverkusen,  
Germany.

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

### **België/Belgique/Belgien**

Bayer SA-NV  
Tél/Tel: +32-(0)2-535 63 11

### **България**

Байер България ЕООД  
Тел.: +359-(0)2 81 401 01

### **Česká republika**

Bayer s.r.o.  
Tel: +420 271 730 661

### **Danmark**

Bayer A/S  
Tlf: +45-45-235 000

### **Deutschland**

Bayer Vital GmbH  
Tel: +49-(0)214-30 513 48

### **Eesti**

Bayer OÜ  
Tel: +372 655 85 65

### **Ελλάδα**

Bayer Ελλάς ABEE  
Τηλ: +30 210 618 75 00

### **España**

Química Farmacéutica Bayer S.L.  
Tel: +34-93-495 65 00

### **France**

Bayer Santé  
Tél: +33-(0)3-28 16 34 00

### **Ireland**

Bayer Limited  
Tel: +353-1-299 93 13

### **Ísland**

Icepharma hf.  
Sími: +354-540 8000

### **Italia**

Bayer S.p.A.  
Tel: +39-02-397 81

### **Κύπρος**

NOVAGEM Limited  
Τηλ: +357-22-747 747

### **Luxembourg/Luxemburg**

Bayer SA-NV  
Tél/Tel: +32-(0)2-535 63 11

### **Magyarország**

Bayer Hungária Kft.  
Tel.: +36-1-487-4100

### **Malta**

Alfred Gera and Sons Ltd.  
Tel: +356-21 44 62 05

### **Nederland**

Bayer B.V., Bayer Schering Pharma  
Tel: +31-(0)297-28 06 66

### **Norge**

Bayer AS  
Tlf:+47-24 11 18 00

### **Österreich**

Bayer Austria Ges. m. b. H.  
Tel: +43-(0)1-711 460

### **Polska**

Bayer Sp. z o.o.  
Tel.: +48-22-572 35 00

### **Portugal**

Bayer Portugal S.A.  
Tel: +351-21-416 42 00

### **România**

SC Bayer SRL.  
Tel: +40-21-528 59 00

### **Slovenija**

Bayer d. o. o.  
Tel: +386-1-58 14 400

### **Slovenská republika**

Bayer, spol. s r.o.  
Tel: +421 2-59 21 31 11

### **Suomi/Finland**

Bayer Oy, Bayer Schering Pharma  
Puh/Tel: +358-20-785 21

### **Sverige**

Bayer AB  
Tel: +46-(0)8-580 223 00

**Latvija**

SIA Bayer

Tel: +371 67 845 563

**Lietuva**

UAB Bayer

Tel: +370 5 23 36 868

**United Kingdom**

Bayer plc

Tel: +44-(0)1635-56 30 00

**This leaflet was last approved in 09/2010**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>